

Cannabinoids and the Urinary Bladder

Evangelia Bakali^{1,2*} and Douglas G Tincello^{1,2}¹University Hospitals of Leicester, NHS trust, Leicester, UK²Reproductive Sciences Section, Department of Cancer Studies and Molecular Medicine, University of Leicester, UK

Abstract

The presence of the Endocannabinoid System (ECS) in the urinary bladder has led to speculation that endocannabinoid-signalling is involved in the signal transduction pathways regulating bladder relaxation and may be involved in pathophysiological processes of the bladder. On the basis of this evidence, it was postulated that the binding of endocannabinoids to the cannabinoid receptors (CB₁ and CB₂) may result in relaxation of the urinary bladder during the filling phase. Dysregulation of the ECS in human bladder may be responsible for the aetiopathogenesis of Overactive Bladder Syndrome (OAB) and Detrusor Overactivity (DO).

Keywords: Cannabinoids; Endocannabinoids; Cannabinoid receptors; Endocannabinoid system

Introduction

Over the past decade, interest in the role of endocannabinoids in regulating many mammalian processes has increased and has been proposed to be involved in the signal transduction mechanism regulating micturition [1,2]. In a sub-analysis of a multicentre, randomized controlled trial of Cannabis in Multiple Sclerosis (CAMS) the effect of cannabinoids on reducing urge incontinence episodes without affecting voiding in patients with multiple sclerosis and Neurogenic Detrusor Overactivity (NDO) was tested [3]. 630 patients were randomized to receive an oral administration of the cannabis extract, Δ^9 -Tetrahydrocannabinol (THC) or matched placebo. Based on incontinence diaries there was a 25% reduction ($p=0.005$) in the cannabis extract group and THC showed a 19% reduction ($p=0.039$) in urinary incontinence episodes relative to placebo suggesting cannabis may modulate detrusor function [3]. This clinical effect of cannabis is supported by the localization and increased density of suburothelial CB₁ nerve fibres in patients with idiopathic detrusor overactivity and painful bladder syndrome compared with controls ($p=0.0123$ and $p=0.0013$ respectively) [2]. However, there are several possible CB receptor isoforms and subtypes and their anatomical distribution, through which the Δ^9 -THC effect is mediated, remains unknown. Since Δ^9 -THC acts on the brain, improvement in urgency and urinary incontinence episodes observed in the CAMS study might be attributed to the effects of Δ^9 -THC at any point in the peripheral nervous system and/or in the micturition centres of the central nervous system.

Historical Review

Cannabis consists of the aerial, seeds and root parts of Cannabis sativa, which is an annual herb indigenous to central and western Asia and is cultivated in other tropical and temperate regions for the fibre used to produce ropes and carpets [4]. There have been more than 60 cannabinoids identified in Cannabis extracts of which the most abundant compound which induces the majority of the psychotropic effects of cannabis, is Δ^9 -THC [5]. Other constituents include cannabiniol, cannabidiol, cannabigerol, cannabichromene and the relative acids [5]. Cannabis has been mentioned in early Hindu and Chinese medicine and its use spread through Persia to Arabia at around the time of the 10th century [6]. The therapeutic effects of cannabinoids were studied in the early 19th Century Irish physician Sir William B. O'Shaughnessy, who demonstrated the potential treatment in a range of disorders including cholera, rheumatic diseases, delirium and infantile convulsions [7]. Historically cannabis has been used in obstetrics and gynaecology for the treatment of menstrual irregularity, dysmenorrhoea, hyperemesis gravidarum, childbirth, postpartum

haemorrhage, menopausal symptoms and urinary symptoms [8]. More common therapeutic applications of cannabis include analgesia, migraine, muscle spasms, seizures, attenuation of nausea and vomiting of cancer chemotherapy, anti-rheumatic and antipyretic actions [8,9].

The pharmacological effects of cannabinoids are mediated by two types of G Protein-Coupled Receptors (GPCR) called CB₁ and CB₂. CB₁ was first identified in 1988 and subsequently cloned from rat cerebral cortex in 1990 [10,11]. It is most widely expressed in central nervous system regions involved with pain transmission and is the most abundant GPCR in the brain [12]. It has also been located in a considerably lower concentration on neurons of peripheral tissues including the heart, vas deferens, urinary bladder and small intestine [12]. The CB₂ receptor was cloned from human promyelocytic leukaemia cells (HL-60 cells) in 1993 and is mainly expressed in immune tissues but is also expressed in low levels in the CNS in both microglia and some neurons [13,14]. The localization of CB₂ receptors in immune tissues implies that some cannabinoid-induced immunosuppression involves a receptor-mediated process. The cannabinoid receptors are activated by natural ligands with arachidonyl ethanolamine (anandamide) being the first endogenous ligand to be isolated. Anandamide mimics the effects of Δ^9 -THC by binding to CB receptors, but lacks the psychocactive effects probably because it is highly susceptible to enzymatic hydrolysis [10,15].

The Endocannabinoid System

The Endocannabinoid System (ECS) consists of the cannabinoid receptors, the endogenous ligands for the cannabinoid receptors, the enzymes involved in the synthesis and degradation of these ligands and the transport systems involved in the transfer of these ligands across the cell membrane.

Cannabinoid Receptors

There are currently three known cannabinoid receptors; CB₁, CB₂, G protein-coupled receptor 55 (GPR55), which are GPCRs activated

***Corresponding author:** Dr. Evangelia Bakali, Reproductive Sciences Section, Department of Cancer Studies and Molecular Medicine, University of Leicester, Robert Kilpatrick Clinical Sciences Building, P.O. Box 65, Leicester LE2 7LX, UK, Tel: +44 116 252 3165; E-mail: eb59@le.ac.uk

Received July 17, 2013; Accepted August 13, 2013; Published August 16, 2013

Citation: Bakali E, Tincello DG (2013) Cannabinoids and the Urinary Bladder. Gynecol Obstet 3: 163. doi:10.4172/2161-0932.1000163

Copyright: © 2013 Bakali E, et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

by endocannabinoid ligands that are arachidonic acid-derived lipid mediators [16]. There are two principal signal transduction pathways involving the cannabinoid receptors; the Cyclic-adenosine monophosphate (cAMP) signal pathway and the phosphatidylinositol signal pathway, which are mediated by the various subunits of G-proteins [16]. Most GPCRs are capable of activating more than one G_{α} -subtype, but they show a preference for one subtype over another [16]. The effector of both the $G_{\alpha s}$ and $G_{\alpha i/o}$ pathways is the enzyme Adenylate Cyclase (AC), which catalyzes the conversion of Cytosolic Adenosine triphosphate (ATP) to cAMP [17]. This mechanism is stimulated by G-proteins of the $G_{\alpha s}$ class and conversely, interaction with G_{α} subunits of the $G_{\alpha i/o}$ type inhibits AC from generating cAMP. [17] The effector of the $G_{\alpha q/11}$ pathway is phospholipase C- β (PLC β), which catalyzes the cleavage of membrane-bound phosphatidylinositol 4,5-bisphosphate (PIP₂) into the second messengers inositol (1,4,5) trisphosphate (IP₃) and Diacylglycerol (DAG)[17]. IP₃ acts on IP₃ receptors found in the membrane of the Endoplasmic Reticulum (ER) to elicit Ca²⁺ release from the ER, while DAG diffuses along the plasma membrane where it may activate any membrane localized forms of a second ser/thr kinase called Protein Kinase C (PKC) [17]. Since many isoforms of PKC are also activated by increases in intracellular Ca²⁺, both these pathways can also converge on each other to signal through the same secondary effector [18]. Elevated [Ca²⁺]_i also binds and allosterically activates proteins called calmodulins, which in turn go on to bind and activate enzymes such as the Ca²⁺/calmodulin-dependant kinases (CAMKs) [19]. Finally, the effectors of the $G_{\alpha 12/13}$ pathway are three RhoGEFs (p115-RhoGEF, PDZ-RhoGEF, and LARG), which, when bound to $G_{\alpha 12/13}$ allosterically activate the cytosolic small GTPase, Rho [19]. Once bound to GTP, Rho can then go on to activate various proteins responsible for cytoskeleton regulation such as Rho-kinase (ROCK) [19]. Most GPCRs that couple to $G_{\alpha 12/13}$ also couple to other sub-classes, often $G_{\alpha q/11}$.

Endocannabinoids



After the cannabinoid receptors were identified as the molecular targets for Δ^9 -THC, natural compounds, which bind to these receptors, were discovered. This group of bioactive lipid signalling molecules was collectively referred to as endogenous cannabinoids or endocannabinoids. *N*-arachidonylolethanolamide (anandamide, AEA) was the first endogenous ligand identified for the cannabinoid receptors in 1992, following its isolation from porcine brain [20]. Since then, a number of bioactive lipid signalling molecules with differing affinities for the cannabinoid receptors have been identified. Additional endocannabinoids include, *N*-docosatetra-7,10,13,16-enylethanolamine, 2-arachidonylglycerol (2-AG), 2-arachidonylglyceryl ether (noladin ether), *O*-arachidonoyl ethanolamine (virodhamine), *N*-dihomo- γ -linoenoyl ethanolamine, *N*-docosatetraenoyl ethanolamine, oleamide, *N*-Arachidonoyl Dopamine (NADA) and *N*-Oleoyl Dopamine (OLDA) (Figure 1). Potency determinations are complicated by the possibility of differential susceptibility of endogenous ligands to enzymatic conversion.

Biosynthesis and degradation of *N*-acylethanolamides

AEA synthesis involves a series of enzymatic reactions, the final stage of which involves the enzyme *N*-arachidonoylphosphatidylethanolamine specific phospholipase D (NAPE-PLD). NAPE-PLD can be stimulated by Ca²⁺, Mg²⁺, Co²⁺, Mn²⁺, Ba²⁺ and Sr²⁺ and other organic cations [21]. Whilst spermine, spermidine, and putrescine are also stimulatory [21]. Initial characterization of NAPE-PLD revealed the enzyme to be membrane associated and it lacks the ability to catalyze a transphosphatidyl transfer reaction, which is a common feature of known

PLDs [22]. NAPE-PLD is the first PLD-type phosphodiesterase which belongs to the metallo- β -lactamase family [23]. Unlike classical neurotransmitters and neuropeptides, its primary product, AEA is not stored in vesicles but synthesized and released "on demand" in response to physiological and pathological stimuli, hormones neurotransmitters and depolarizing agents from its direct biosynthetic precursor *N*-arachidonoylphosphatidylethanolamine (NAPE) a phospholipid commonly found in biological membranes [24,25]. Figure 2 shows an outline of the major pathways through which anandamide and 2-AG are produced and degraded.

Fatty Acid Amide Hydrolase (FAAH) is the enzyme primarily involved in the hydrolysis of AEA, but can also degrade other endocannabinoids. FAAH was first cloned and purified from rat liver



microsomes but is present in many other tissues and often in tissues containing CB₁ and CB₂ receptors [26]. In addition to FAAH, AEA can also be degraded by Palmitoylethanolamide-Preferring Acid Amidase (PAA), cyclooxygenase-2, lipoxygenases and cytochrome P450 [27]. 2-AG is the second member of the endogenous cannabinoid family to be identified, which binds to both CB₁ and CB₂ receptors with similar affinities to AEA, although 2-AG has a higher affinity for CB₂ receptors than CB₁ [12]. The synthesis of 2-AG depends on the conversion of 2-arachidonate-containing phosphoinositides to diacylglycerols and their subsequent transformation to 2-arachidonylglycerol by action of two Diacylglycerol Lipase (DAGL) isozymes, DAGLa and DAGLβ. Following their synthesis and release, these endocannabinoids are removed from their sites of action by cellular uptake and degraded by their enzymes. 2-AG is mainly degraded by Monoacylglycerol Lipase (MAGL) but a small amount is also degraded by FAAH.

Synthetic Ligands

Cannabinoid agonists are classified by chemical structure into four main groups: classical; non-classical; aminoalkylindoles; and eicosanoids [28] (Figure 3). Classical cannabinoids are dibenzopyrane derivatives and include Δ⁹-THC, while non-classical consists of a bicyclic and tricyclic analogue of Δ⁹-THC that lacks a pyran ring [28]. One major practical difficulty associated with cannabinoid research both in vivo and in vitro, is the high lipophilicity and low water solubility of most CB₁ and CB₂ receptor ligands as this necessitates the use of a non-aqueous vehicle such as ethanol, Dimethyl Sulphoxide (DMSO), polyvinylpyrrolidone, Tween 80, Cremophor, Emulphor, bovine serum albumin, or water soluble emulsion Tocrisolve 100, which is a mixture of soya oil, Pluronic F68 and water to get the compound of interest to the cell surface [29]. It also means that these compounds “stick” to equipment during treatment, which needs to be taken into consideration during experimental procedures.

Cannabinoid receptor signaling


Calcium acts as an intracellular messenger where it plays a key role in regulating basic cellular responses, such as migration and proliferation [30]. Under resting conditions, cytoplasmic calcium concentration is maintained at approximately 100nM [30]. When stimulated, calcium enters the cell from extracellular stores via ion channels in the plasma membrane or it is released from intracellular stores through channels and receptors in the endoplasmic reticulum [30]. These channels may be activated and modulated by second messengers including IP₃, which is produced by binding of ligands, such as ATP, to GPCRs [31].

The CB₁ receptor is a member of the rhodopsin subfamily of GPCRs [32]. There are three cytosolic loops and a putative fourth loop formed by palmitoylation at the juxtamembrane C-terminal region, which contribute to the activation of the G-proteins [33]. The proximal CB₁ receptor intracellular C-terminal domain is critical for G-protein coupling and the distal C-terminal tail domain modulates signal transduction [33]. Most cannabinoid effects are sensitive to Pertussis Toxin (PTX) implicating a CB₁ and CB₂ receptor coupling to a G_{i/o} protein [34]. The binding of endocannabinoids and cannabinoids to CB₁ and CB₂ results in a decrease of intracellular cAMP levels and activation of mitogen-activated protein kinase through the coupled G_{i/o} proteins [34-36]. Cannabinoid-mediated inhibition of cAMP has been demonstrated in slices of rat hippocampus, striatum, cerebral cortex and cerebellum [36]. CB₁ can also stimulate the formation of cAMP through G_s under certain conditions [37]. It may also be that CB₁ receptors can exist as two distinct subpopulations, one coupled to G_{i/o} proteins and the other to G_s [38,39]. The level of cytosolic cAMP may then determine the activity of various ion channels as well as members of the ser/thr specific protein kinase A (PKA) family [32,40,41]. Thus cAMP is considered a second messenger and PKA a secondary effector.

In addition, activation of CB₁ receptor modulates ion channels through G_{i/o} proteins leading to the activation of A-type and inwardly rectifying potassium channels [42-45]. This is due to decreased phosphorylation of the channels, as protein kinase A activity is decreased due to cannabinoid induced inhibition of AC [45]. Thus cannabinoids increase the efflux of potassium. In addition, activation of CB₁ causes a cAMP-independent, but G_{i/o}-dependent inhibition of N-type and P/Q-type calcium channels and activation of inwardly rectifying potassium channel proteins (e.g. GIRK1, GIRK2), leading to a decrease calcium influx and increase in potassium efflux [42-44].

Similarly, CB₁, CB₂ receptors can modulate AC and MAP kinase activity, through their ability to couple to G_{i/o} proteins [46]. The MAP kinase pathway is a key signalling mechanism that regulates many cellular functions such as cell growth, transformation, differentiation, gene expression and apoptosis [47]. Activation of the MAP kinase pathway is associated with the activation of a tyrosine kinase-linked receptor which activates the intracellular G protein Ras and sets up a signaling cascade beginning with the activation of the serine/threonine kinase Raf (MAP kinase kinase kinase) [32]. Raf activates MAP kinase kinase (MEK) leading to phosphorylation and activation of MAP kinase, which can phosphorylate various cytoplasmic and nuclear proteins [32]. CB₁ receptors have been shown to link positively to MAP kinase [48]. However, in contrast to CB₁, CB₂ receptor stimulation is believed not to modulate ion channel function as seen in AtT-20 cells transfected with CB₂ receptors and Xenopus oocytes transfected with CB₂ [49,50]. In addition, unlike CB₁ receptors, CB₂ receptors do not appear to couple to G_s, suggesting there is a difference between CB₁ and CB₂ receptor signalling [51].

There is evidence that GPR55 is a novel cannabinoid receptor that has a different signalling pathway to that of CB₁ and CB₂ [52,53]. GPR55 is also a rhodopsin-like GPCR, which has been implicated in diverse physiological and pathological processes such as inflammatory and neuropathic pain, bone development and cancer. However, GPR55 shares only low amino acid sequence identity with CB₁ (13.5%) and CB₂ (14.4%) and lacks the typical functional response elicited by these receptors [54]. Activation of the GPR55 receptor coupled to the G_q, G₁₂, RHoA, actin, phospholipase C pathway triggers the release of Ca²⁺ from IP₃R-gated stores, which leads to increased intracellular Ca²⁺ [53] (Figure 4). GPR55 can be activated by Lysophosphatidylinositol (LPI), which is an agonist, which can be antagonized by CP55940 and cannabidiol.



Cannabis and the Urinary Bladder

Cannabinoid Receptor distribution in the urinary bladder

The effect of cannabis on DO symptoms is probably mediated through a mechanism that depends on endocannabinoids [3]. The mechanism of this effect is far from clear and published data on the expression and functional sites of cannabinoid receptors in the bladder are contradictory. It is thought that endocannabinoids bind to CB₁ and CB₂, resulting in relaxation of the detrusor muscle during the filling phase [55,56]. CB₁ receptors are mainly found at the central and peripheral neuron terminals of the bladder, inhibiting neurotransmitter release [55]. Several studies have localized both cannabinoid receptors in the urinary bladder of humans rats mice and monkeys [2,55-60]. The localization of CB₁ receptors has been described to be in the urothelium and nerve fibres of the suburothelium and in human and rat detrusor muscle [2,58,60]. However, another study did not detect the CB₁ receptor in rat urothelium or nerve fibres but reported immunoreactivity for CB₂ in these structures and in ganglion cells of the outflow region [1,55]. In addition, human bladder studies identifying the presence of gene transcripts by quantitative Polymerase Chain Reaction (qPCR) and tissue expression and localization by Immunohistochemistry (IHC), revealed a higher abundance of the CB₁ receptor in the urothelium compared to the detrusor [57]. Similar results were found for CB₂ but overall, receptor protein expression was much lower when compared to CB₁ receptor protein expression [57].

Cannabinoid Receptor function in the urinary bladder

Studies have demonstrated that the activation of presynaptic CB₁ and CB₂ receptors inhibit electrically evoked contractions in isolated mammalian tissue when using THC and the non-selective CB receptor agonists CP55940, CP55244, JWH015, which corresponds to the localization of CB₁ receptors in nerve fibres of the detrusor muscle [55,60-62]. In isolated mouse bladder, several cannabinoid receptor agonists, including WIN 55212-2, Δ⁹-THC and anandamide, inhibited electrically-evoked bladder contractions in a concentration dependent manner [61]. In the same study, it was shown that the inhibitory effect was not a post-synaptic effect since contractile responses to muscarinic or purinergic receptor agonists were unaffected by pre-treatment with Δ⁹-THC [61]. In rat detrusor muscle, cannabitor (a CB₂ selective agonist) did not have any effects on nerve-induced contractions [1]. Similarly, in a study where human bladder muscle strips were used, there was no inhibitory effect of the non-selective CB agonist, WIN 55212-2, on Electrical Field Stimulation (EFS) evoked contraction [62].

In contrast, another study found an attenuation of EFS evoked human detrusor contraction in the presence of both CB₁ (ACEA) and CB₂ (GP1a) agonists [57]. These findings suggest cannabinoids act on pre-junctional nerve endings attenuating contractile responses. These data, however, must be interpreted with caution because quantification of the effect by GP1a or vehicle (dimethyl-sulfoxide) control experiments were not presented [57]. Supporting that cannabinoids act on pre-junctional nerve endings to attenuate a contractile response, Gratzke et al. demonstrated co-localization of vesicular acetylcholine transporter protein (VACHT) nerve structures and CB2 immunoreactive terminal varicosities. They also showed inhibitory effects of CP55, 940 on nerve mediated contractions but not on carbachol induced contractions in detrusor preparations, suggesting a modulatory function of CB₂ on cholinergic neurotransmission [55]. Similarly, cannabitor (a CB₂ agonist) did not attenuate carbachol-induced contractions in isolated rat detrusor tissue, suggesting that the action of the CB₂ receptor is not directly involved in post-junctional regulation of smooth muscle contractility [1]. A recent study showed that both pure Cannabidiol (CBD) and Cannabis Sativa extract enriched with CBD also termed as "CBD Botanic Drug Substance" (CBD BDS), which are devoid of psychotropic activity, inhibited human and rat bladder contractility via a postsynaptic site of action [63].

The differences seen between the results of these studies may be due to inter-species differences in cannabinoid receptor expression and distribution, the effect of these receptors on the release of contractile transmitters and anatomical variations in bladder innervation. Inter-species differences in the neuroanatomy of the mammalian bladders are known to exist [62]. For example there are several parasympathetic ganglia in isolated bladder tissue from guinea pigs and humans while there are none in the urinary bladders of mice and rats [64,65].

Cystometric studies have shown an increase of the micturition threshold in rats receiving systemic cannabinoids in normal and inflamed conditions induced by acetic acid, cyclophosphamide or turpentine oil [66,67]. These effects were stronger when the cannabinoids were administered through a close-arterial route rather than systemically through the tail vein of the rat, supporting the hypothesis of a local regulatory role of the cannabinoid system in the micturition reflex [67]. The mechanism by which cannabinoid receptors could modulate this reflex is by the presence of CB₁ receptors in the afferent nerve fibre endings located in the suburothelial layer, which is supported by in vitro studies where CB₁ agonists reduce neuronal activity and attenuate bladder contractility as a result of electrical field stimulation in isolated mouse bladder strips [61,68]. In rats, anandamide, WIN 55212-2 (synthetic CB non-selective agonist), and Ajulemic acid (IP-751) (synthetic THC analogue), suppress normal bladder activity and the urinary frequency induced by bladder irritation suggesting the inhibitory effects are least in part mediated by CB₁ receptors [66,67,69]. A recent study, showed that CB₂ receptor mediated signals using a high affinity CB₂ receptor selective agonist, cannabitor 3.0 mg/kg, increased the micturition intervals and volumes by 52% (p < 0.05) and 96% (p < 0.01), respectively, and increased threshold and flow pressures by 73% (p < 0.01) and 49% (p < 0.001), respectively, in conscious rats during cystometry [1]. It has not been clarified if these actions are related to CB receptors in the central nervous system, at peripheral sites in the lower urinary tract, or both. Furthermore, it is not known which of the two CB receptor subtypes is mainly responsible for the regulation of micturition in the different species.

Cannabinoid receptors as therapeutic targets

The most studied cannabinoid compound is Cannabidiol (CBD)




Figure 4: GPR55 signalling.

which exerts a number of pharmacologic effects such as analgesic, anti-inflammatory, antioxidant, and anti-tumoral [70]. It has been clinically evaluated for the treatment of anxiety, psychosis, and movement disorders and has been found to have a safe clinical profile [70]. CBD is the main component of Sativex, which also contains Δ^9 -THC, a cannabis-derived drug used for the treatment of pain and spasticity associated with multiple sclerosis. Sativex is licensed for this indication in patients with multiple sclerosis. In a clinical survey, administration of Δ^9 -THC improved nocturia and detrusor overactivity in patients with multiple sclerosis [71]. To date, a small number of open-label and placebo-controlled studies have demonstrated that oral administration of cannabinoids may alleviate OAB/DO symptoms as first line. Most of these studies have been carried out on patients with advanced multiple sclerosis using preparations containing Δ^9 -THC and/or CBD. One such study using Sativex, showed a reduction in urgency, number of incontinence episodes, frequency and nocturia in patients with multiple sclerosis [72,73]. Other cannabinoid receptor agonists are already used clinically to suppress nausea and vomiting provoked by anticancer drugs (nabilone) or to boost the appetite of AIDS patients but these have not been studied for their effects upon urinary symptoms [9].

However, the oral use of cannabinoids may induce undesirable CNS effects including hypoactivity, hypothermia and catalepsy, but may in turn improve OAB symptoms, which are known to be afferently mediated [3,74]. What remains unclear is whether the latter beneficial effects are centrally mediated or whether a local bladder component acting on the afferent bladder pathway, plays a significant role. There are no human data that exists which can answer this question. Data from animal studies support a local effect on bladder afferents where cannabinoid administration systemically and intravesically, improved parameters associated with OAB and DO [55,75].

In addition to using an intravesical route of administration for cannabinoid drugs in order to bypass the CNS effects associated with activation of CB1, the use of CB2 agonists and FAAH inhibitors is being explored and appear promising [76,77]. There is emerging evidence that activation of CB2 inhibits tissue inflammation and has analgesic properties [78-80]. The CB2 subtype is mainly expressed outside the CNS, as described earlier, so it can act as a potential endocannabinoid target where analgesic effects may be separated from psychotropic effects by activating the peripheral receptors. In addition, pharmacological targeting of the homeostasis of endogenous cannabinoids by manipulating the degradation enzymes, may also offer the possibility of avoiding the CNS side effects of exogenous cannabinoids. FAAH, an enzyme that specifically degrades anandamide has been localised in the urinary bladder [56,77,81]. Inhibition of FAAH activity with FAAH inhibitor Oleoyl Ethyl Amide (OEA), significantly increased inter-contraction intervals, micturition volume, bladder capacity and threshold pressure urodynamic parameters in rats which reflect sensory functions of micturition. These effects were prevented by a selective CB2 antagonist. Similarly, another FAAH inhibitor, URB597 has been found to have a functional role in the colon, where FAAH has been localized by reducing inflammation [77,82,83]. The use of a FAAH inhibitor needs to be explored further in the urinary bladder because it may be the way forward in treating OAB symptoms. However, the complexity of the endocannabinoid system at the tissue level may mean that we are still a long way from obtaining a clinically useful compound for treatment.

The Future

Modulation of the endocannabinoid system is currently being investigated for a wide range of potential therapeutic

applications including smoking cessation, treatment of obesity, epilepsy and other CNS related conditions. Similarly, the presence of the endocannabinoid system in the urinary bladder has led to speculation that endocannabinoid-signalling is involved in the signal transduction pathways regulating bladder relaxation and may be involved in pathophysiological processes of the bladder. This role of the endocannabinoids in the lower urinary tract supports their therapeutic potential in conditions of OAB and DO, whereas evidence already exists for their role in bladder inflammation [2,59,75,84]. There are still a number of unanswered questions in the understanding of cannabinoid pharmacology in the urinary bladder. Clearly, further research is required to investigate the role of cannabinoid receptors and their exogenous modulators on bladder control prior to embarking on a clinical trial involving cannabinoids and healthy volunteers with OAB. The inhibitory effects of CB2 and the effect of FAAH inhibitors on lower urinary tract control should be the focus of future studies.

Acknowledgments

This work was funded by Wellbeing of Women Training fellowship, UK.

References

1. Gratzke C, Streng T, Stief CG, Downs TR, Alroy I, et al. (2010) Effects of cannabimol, a novel selective cannabinoid 2 receptor agonist, on bladder function in normal rats. *Eur Urol* 57: 1093-1100.
2. Mukerji G, Yiangou Y, Agarwal SK, Anand P (2010) Increased cannabinoid receptor 1-immunoreactive nerve fibers in overactive and painful bladder disorders and their correlation with symptoms. *Urology* 75: 1514.
3. Freeman RM, Adekanmi O, Waterfield MR, Waterfield AE, Wright D, et al. (2006) The effect of cannabis on urge incontinence in patients with multiple sclerosis: a multicentre, randomised placebo-controlled trial (CAMS-LUTS). *Int Urogynecol J Pelvic Floor Dysfunct* 17: 636-641.
4. Mechoulam R (1986) Chapter 1: the pharmacology of Cannabis sativa Cannabinoids as Therapeutic Agents, Birkhauser, UK.
5. Samuelsson G (1992) Drugs of natural origin: a textbook of pharmacognosy. Swedish Pharmaceutical Press, Denmark.
6. Abel EL (1980) Marijuana, the first twelve thousand years 289. Plenum Press, USA.
7. O'Shaughnessy WB (1843) On the Preparations of the Indian Hemp, or Gunjah: Cannabis Indica Their Effects on the Animal System in Health, and their Utility in the Treatment of Tetanus and other Convulsive Diseases *Prov Med J Retrospect Med Sci* 5: 363-369.
8. Hollister LE (1986) Health aspects of cannabis. *Pharmacol Rev* 38: 1-20.
9. Ben Amar M (2006) Cannabinoids in medicine: A review of their therapeutic potential. *J Ethnopharmacol* 105: 1-25.
10. Devane WA, Dysarz FA 3rd, Johnson MR, Melvin LS, Howlett AC (1988) Determination and characterization of a cannabinoid receptor in rat brain. *Mol Pharmacol* 34: 605-613.
11. Matsuda LA, Lolait SJ, Brownstein MJ, Young AC, Bonner TI (1990) Structure of a cannabinoid receptor and functional expression of the cloned cDNA. *Nature* 346: 561-564.
12. Pertwee RG, Ross RA (2002) Cannabinoid receptors and their ligands. *Prostaglandins Leukot Essent Fatty Acids* 66: 101-121.
13. Munro S, Thomas KL, Abu-Shaar M (1993) Molecular characterization of a peripheral receptor for cannabinoids. *Nature* 365: 61-65.
14. Gong JP, Onaivi ES, Ishiguro H, Liu QR, Tagliaferro PA, et al. (2006) Cannabinoid CB2 receptors: immunohistochemical localization in rat brain. *Brain Res* 1071: 10-23.
15. Deutsch DG, Chin SA (1993) Enzymatic synthesis and degradation of anandamide, a cannabinoid receptor agonist. *Biochem Pharmacol* 46: 791-796.
16. Wang J, Ueda N (2009) Biology of endocannabinoid synthesis system. *Prostaglandins Other Lipid Mediat* 89: 112-119.
17. Lodish H, Berk A, Zipursky SL, Paul Matsudaira, David Baltimore, et al. (2000)

- G Protein- Coupled Receptors and Their Effectors. Molecular Cell Biology (4th Edition), W.H. Freeman Publishers, USA.
18. Massotte D, Kieffer BL (2005) Structure-Function Relationships In: The G Protein-Coupled Receptors Handbook. Humana press, New jersey, USA.
 19. Hwangpo TN, Iyengar R (2005) Heterotrimeric G proteins and their effector pathways In: The G Protein-Coupled Receptors Handbook. Humana press, New jersey, USA.
 20. Devane WA, Hanus L, Breuer A, Pertwee RG, Stevenson LA, et al. (1992) Isolation and structure of a brain constituent that binds to the cannabinoid receptor. *Science* 258: 1946-1949.
 21. Liu Q, Tonai T, Ueda N (2002) Activation of N-acyl ethanolamine-releasing phospholipase D by polyamines. *Chem Phys Lipids* 115: 77-84.
 22. Petersen G, Hansen HS (1999) N-acylphosphatidylethanolamine-hydrolysing phospholipase D lacks the ability to transphosphatidylate. *FEBS Lett* 455: 41-44.
 23. Wang J, Okamoto Y, Morishita J, Tsuboi K, Miyatake A, et al. (2006) Functional analysis of the purified anandamide-generating phospholipase D as a member of the metallo-beta-lactamase family. *J Biol Chem* 281: 12325-12335.
 24. Piomelli D (2003) The molecular logic of endocannabinoid signalling. *Nat Rev Neurosci* 4: 873-884.
 25. De Petrocellis L, Cascio MG, Di Marzo V (2004) The endocannabinoid system: a general view and latest additions. *Br J Pharmacol* 141: 765-774.
 26. Cravatt BF, Giang DK, Mayfield SP, Boger DL, Lerner RA, et al. (1996) Molecular characterization of an enzyme that degrades neuromodulatory fatty-acid amides. *Nature* 384: 83-87.
 27. Pertwee RG (2005) Pharmacological actions of cannabinoids. *Handb Exp Pharmacol* : 1-51.
 28. Nocerino E, Amato M, Izzo AA (2000) Cannabis and cannabinoid receptors. *Fitoterapia* 71 Suppl 1: S6-12.
 29. Howlett AC, Barth F, Bonner TI, Cabral G, Casellas P, et al. (2002) International Union of Pharmacology. XXVII. Classification of cannabinoid receptors. *Pharmacol Rev* 54: 161-202.
 30. Shabir S, Southgate J (2008) Calcium signalling in wound-responsive normal human urothelial cell monolayers. *Cell Calcium* 44: 453-464.
 31. Galione A, Churchill GC (2002) Interactions between calcium release pathways: multiple messengers and multiple stores. *Cell Calcium* 32: 343-354.
 32. Demuth DG, Molleman A (2006) Cannabinoid signalling. *Life Sci* 78: 549-563.
 33. Nie J, Lewis DL (2001) The proximal and distal C-terminal tail domains of the CB1 cannabinoid receptor mediate G protein coupling. *Neuroscience* 107: 161-167.
 34. Howlett AC, Qualy JM, Khachatrian LL (1986) Involvement of Gi in the inhibition of adenylate cyclase by cannabinimetic drugs. *Mol Pharmacol* 29: 307-313.
 35. Howlett AC, Fleming RM (1984) Cannabinoid inhibition of adenylate cyclase. Pharmacology of the response in neuroblastoma cell membranes. *Mol Pharmacol* 26: 532-538.
 36. Bidaut-Russell M, Devane WA, Howlett AC (1990) Cannabinoid receptors and modulation of cyclic AMP accumulation in the rat brain. *J Neurochem* 55: 21-26.
 37. Howlett AC (2005) Cannabinoid receptor signaling. *Handb Exp Pharmacol* : 53-79.
 38. Calandra B, Portier M, Kernéis A, Delpech M, Carillon C, et al. (1999) Dual intracellular signaling pathways mediated by the human cannabinoid CB1 receptor. *Eur J Pharmacol* 374: 445-455.
 39. Bonhaus DW, Chang LK, Kwan J, Martin GR (1998) Dual activation and inhibition of adenylate cyclase by cannabinoid receptor agonists: evidence for agonist-specific trafficking of intracellular responses. *J Pharmacol Exp Ther* 287: 884-888.
 40. Mu J, Zhuang SY, Kirby MT, Hampson RE, Deadwyler SA (1999) Cannabinoid receptors differentially modulate potassium A and D currents in hippocampal neurons in culture. *J Pharmacol Exp Ther* 291: 893-902.
 41. Mu J, Zhuang SY, Hampson RE, Deadwyler SA (2000) Protein kinase-dependent phosphorylation and cannabinoid receptor modulation of potassium A current (IA) in cultured rat hippocampal neurons. *Pflugers Arch* 439: 541-546.
 42. Mackie K, Hille B (1992) Cannabinoids inhibit N-type calcium channels in neuroblastoma-glioma cells. *Proc Natl Acad Sci U S A* 89: 3825-3829.
 43. Mackie K, Lai Y, Westenbroek R, Mitchell R (1995) Cannabinoids activate an inwardly rectifying potassium conductance and inhibit Q-type calcium currents in AT20 cells transfected with rat brain cannabinoid receptor. *J Neurosci* 15: 6552-6561.
 44. Henry DJ, Chavkin C (1995) Activation of inwardly rectifying potassium channels (GIRK1) by co-expressed rat brain cannabinoid receptors in *Xenopus* oocytes. *Neurosci Lett* 186: 91-94.
 45. Deadwyler SA, Hampson RE, Mu J, Whyte A, Childers S (1995) Cannabinoids modulate voltage sensitive potassium A-current in hippocampal neurons via a cAMP-dependent process. *J Pharmacol Exp Ther* 273: 734-743.
 46. Kobayashi Y, Arai S, Waku K, Sugiura T (2001) Activation by 2-arachidonoylglycerol, an endogenous cannabinoid receptor ligand, of p42/44 mitogen-activated protein kinase in HL-60 cells. *J Biochem* 129: 665-669.
 47. Pearson G, Robinson F, Beers Gibson T, Xu BE, Karandikar M, et al. (2001) Mitogen-activated protein (MAP) kinase pathways: regulation and physiological functions. *Endocr Rev* 22: 153-183.
 48. Galve-Roperh I, Rueda D, Gómez del Pulgar T, Velasco G, Guzmán M (2002) Mechanism of extracellular signal-regulated kinase activation by the CB(1) cannabinoid receptor. *Mol Pharmacol* 62: 1385-1392.
 49. Felder CC, Joyce KE, Briley EM, Mansouri J, Mackie K, et al. (1995) Comparison of the pharmacology and signal transduction of the human cannabinoid CB1 and CB2 receptors. *Mol Pharmacol* 48: 443-450.
 50. McAllister SD, Griffin G, Satin LS, Abood ME (1999) Cannabinoid receptors can activate and inhibit G protein-coupled inwardly rectifying potassium channels in a *Xenopus* oocyte expression system. *J Pharmacol Exp Ther* 291: 618-626.
 51. Glass M, Felder CC (1997) Concurrent stimulation of cannabinoid CB1 and dopamine D2 receptors augments cAMP accumulation in striatal neurons: evidence for a Gs linkage to the CB1 receptor. *J Neurosci* 17: 5327-5333.
 52. Mackie K, Stella N (2006) Cannabinoid receptors and endocannabinoids: evidence for new players. *AAPS J* 8: E298-306.
 53. Lauckner JE, Jensen JB, Chen HY, Lu HC, Hille B, et al. (2008) GPR55 is a cannabinoid receptor that increases intracellular calcium and inhibits M current. *Proc Natl Acad Sci U S A* 105: 2699-2704.
 54. Pertwee RG (2007) GPR55: a new member of the cannabinoid receptor clan? *Br J Pharmacol* 152: 984-986.
 55. Gratzke C, Streng T, Park A, Christ G, Stief CG, et al. (2009) Distribution and function of cannabinoid receptors 1 and 2 in the rat, monkey and human bladder. *J Urol* 181: 1939-1948.
 56. Bakali E, Elliott RA, Taylor AH, Willets J, Konje JC, et al. (2013) Distribution and function of the endocannabinoid system in the rat and human bladder. *Int Urogynecol J* 24: 855-863.
 57. Tyagi V, Philips BJ, Su R, Smaldone MC, Erickson VL, et al. (2009) Differential expression of functional cannabinoid receptors in human bladder detrusor and urothelium. *J Urol* 181: 1932-1938.
 58. Hayn MH, Ballesteros I, de Miguel F, Coyle CH, Tyagi S, et al. (2008) Functional and immunohistochemical characterization of CB1 and CB2 receptors in rat bladder. *Urology* 72: 1174-1178.
 59. Merriam FV, Wang ZY, Guerios SD, Bjorling DE (2008) Cannabinoid receptor 2 is increased in acutely and chronically inflamed bladder of rats. *Neurosci Lett* 445: 130-134.
 60. Walczak JS, Price TJ, Cervero F (2009) Cannabinoid CB1 receptors are expressed in the mouse urinary bladder and their activation modulates afferent bladder activity. *Neuroscience* 159: 1154-1163.
 61. Pertwee RG, Fernando SR (1996) Evidence for the presence of cannabinoid CB1 receptors in mouse urinary bladder. *Br J Pharmacol* 118: 2053-2058.
 62. Martin RS, Luong LA, Welsh NJ, Eglen RM, Martin GR, et al. (2000) Effects of cannabinoid receptor agonists on neuronally-evoked contractions of urinary bladder tissues isolated from rat, mouse, pig, dog, monkey and human. *Br J Pharmacol* 129: 1707-1715.
 63. Capasso R, Aviello G, Borrelli F, Romano B, Ferro M, et al. (2011) Inhibitory effect of standardized cannabis sativa extract and its ingredient cannabidiol on rat and human bladder contractility. *Urology* 77: 1006.

64. Gabella G (1990) Intramural neurons in the urinary bladder of the guinea-pig. *Cell Tissue Res* 261: 231-237.
65. Gilpin CJ, Dixon JS, Gilpin SA, Gosling JA (1983) The fine structure of autonomic neurons in the wall of the human urinary bladder. *J Anat* 137: 705-713.
66. Hiragata S, Ogawa T, Hayashi Y, Tyagi P, Seki S, et al. (2007) Effects of IP-751, ajulemic acid, on bladder overactivity induced by bladder irritation in rats. *Urology* 70: 202-208.
67. Dmitrieva N, Berkley KJ (2002) Contrasting effects of WIN 55212-2 on motility of the rat bladder and uterus. *J Neurosci* 22: 7147-7153.
68. Di Marzo V, Bifulco M, De Petrocellis L (2004) The endocannabinoid system and its therapeutic exploitation. *Nat Rev Drug Discov* 3: 771-784.
69. Jaggar SI, Hasnie FS, Sellaturay S, Rice AS (1998) The anti-hyperalgesic actions of the cannabinoid anandamide and the putative CB2 receptor agonist palmitoylethanolamide in visceral and somatic inflammatory pain. *Pain* 76: 189-199.
70. Izzo AA, Borrelli F, Capasso R, Di Marzo V, Mechoulam R (2009) Non-psychoactive plant cannabinoids: new therapeutic opportunities from an ancient herb. *Trends Pharmacol Sci* 30: 515-527.
71. Consroe P, Musty R, Rein J, Tillery W, Pertwee R (1997) The perceived effects of smoked cannabis on patients with multiple sclerosis. *Eur Neurol* 38: 44-48.
72. Wade DT, Makela P, Robson P, House H, Bateman C (2004) Do cannabis-based medicinal extracts have general or specific effects on symptoms in multiple sclerosis? A double-blind, randomized, placebo-controlled study on 160 patients. *Mult Scler* 10: 434-441.
73. Brady CM, DasGupta R, Dalton C, Wiseman OJ, Berkley KJ, et al. (2004) An open-label pilot study of cannabis-based extracts for bladder dysfunction in advanced multiple sclerosis. *Mult Scler* 10: 425-433.
74. Brady CM, Apostolidis AN, Harper M, Yiangou Y, Beckett A, et al. (2004) Parallel changes in bladder suburothelial vanilloid receptor TRPV1 and neuronal marker PGP9.5 immunoreactivity in patients with neurogenic detrusor overactivity after intravesical resiniferatoxin treatment. *BJU Int* 93: 770-776.
75. Walczak JS, Cervero F (2011) Local activation of cannabinoid CB₁ receptors in the urinary bladder reduces the inflammation-induced sensitization of bladder afferents. *Mol Pain* 7: 31.
76. Guindon J, Hohmann AG (2008) Cannabinoid CB2 receptors: a therapeutic target for the treatment of inflammatory and neuropathic pain. *Br J Pharmacol* 153: 319-334.
77. Strittmatter F, Gandaglia G, Benigni F, Bettiga A, Rigatti P, et al. (2012) Expression of fatty acid amide hydrolase (FAAH) in human, mouse, and rat urinary bladder and effects of FAAH inhibition on bladder function in awake rats. *Eur Urol* 61: 98-106.
78. Barutta F, Piscitelli F, Pinach S, Bruno G, Gambino R, et al. (2011) Protective role of cannabinoid receptor type 2 in a mouse model of diabetic nephropathy. *Diabetes* 60: 2386-2396.
79. Quartilho A, Mata HP, Ibrahim MM, Vanderah TW, Porreca F, et al. (2003) Inhibition of inflammatory hyperalgesia by activation of peripheral CB2 cannabinoid receptors. *Anesthesiology* 99: 955-960.
80. Palazuelos J, Aguado T, Pazos MR, Julien B, Carrasco C, et al. (2009) Microglial CB2 cannabinoid receptors are neuroprotective in Huntington's disease excitotoxicity. *Brain* 132: 3152-3164.
81. Merriam FV, Wang ZY, Hillard CJ, Stuhr KL, Bjorling DE (2011) Inhibition of fatty acid amide hydrolase suppresses referred hyperalgesia induced by bladder inflammation. *BJU Int* 108: 1145-1149.
82. Marqu ez L, Su arez J, Iglesias M, Bermudez-Silva FJ, Rodr guez de Fonseca F, et al. (2009) Ulcerative colitis induces changes on the expression of the endocannabinoid system in the human colonic tissue. *PLoS One* 4: e6893.
83. Storr MA, Keenan CM, Emmerdinger D, Zhang H, Y ce B, et al. (2008) Targeting endocannabinoid degradation protects against experimental colitis in mice: involvement of CB1 and CB2 receptors. *J Mol Med (Berl)* 86: 925-936.
84. Szallasi A, Fowler CJ (2002) After a decade of intravesical vanilloid therapy: still more questions than answers. *Lancet Neurol* 1: 167-172.

Citation: Bakali E, Tincello DG (2013) Cannabinoids and the Urinary Bladder. *Gynecol Obstet* 3: 163. doi:10.4172/2161-0932.1000163

Submit your next manuscript and get advantages of OMICS Group submissions

Unique features:

- User friendly/feasible website-translation of your paper to 50 world's leading languages
- Audio Version of published paper
- Digital articles to share and explore

Special features:

- 250 Open Access Journals
- 20,000 editorial team
- 21 days rapid review process
- Quality and quick editorial, review and publication processing
- Indexing at PubMed (partial), Scopus, EBSCO, Index Copernicus and Google Scholar etc
- Sharing Option: Social Networking Enabled
- Authors, Reviewers and Editors rewarded with online Scientific Credits
- Better discount for your subsequent articles

Submit your manuscript at: <http://www.omicsonline.org/submission/>

